

LIPSCHITZ SPACES AND CALDERÓN-ZYGMUND OPERATORS ASSOCIATED TO NON-DOUBLING MEASURES

JOSÉ GARCÍA-CUERVA AND A. EDUARDO GATTO

ABSTRACT. In the setting of $\mathbb{R}^d$ with an n -dimensional measure μ , we give several characterizations of Lipschitz spaces in terms of mean oscillations involving μ . We also show that Lipschitz spaces are preserved by those Calderón-Zygmund operators T associated to the measure μ for which $T(1)$ is the Lipschitz class 0.

1. INTRODUCTION

In the sequel $\mathbb{R}^d$ will denote the euclidean space of vectors $x = (x_1, \dots, x_d)$ with d real components and μ a Radon n -dimensional measure on $\mathbb{R}^d$. We say that μ is an n -dimensional measure, $n \in \mathbb{R}$, $0 < n \leq d$, if it satisfies the following growth condition: There is a constant C such that

$$(1.1) \quad \mu(B(x, r)) \leq Cr^n$$

for every ball $B(x, r)$ with center $x \in \mathbb{R}^d$ and radius $r > 0$. This allows, in particular, non-doubling measures.

The study of Calderón-Zygmund operators associated to an n -dimensional measure was carried out, in the Lebesgue spaces, by Nazarov, Treil and Volberg (see [NTV1, NTV2]) and also by Tolsa (see [To1, To2]). Further results, dealing with BMO and H^1 and providing boundedness criteria in the spirit of the $T(1)$ or $T(b)$ theorems, were obtained as well (see [NTV3, MMNO, To3])

Date: February 1, 2008.

1991 *Mathematics Subject Classification.* 2000 revision, 42B20, 26B35, 47B38, 47G10.

Key words and phrases. Calderón-Zygmund theory, singular integrals, Lipschitz spaces, BMO, non-doubling measures.

Supported in part by DGES, Spain, under grant BFM2001-0189. It is a pleasure for the second author, to thank the members of the Mathematics Department of Universidad Autónoma de Madrid for their friendly hospitality.

In [GG] we have also studied the theory of fractional integral operators associated to an n -dimensional measure μ on Lebesgue spaces and Lipschitz spaces.

The present paper is devoted to further study the Lipschitz spaces associated to an n -dimensional measure μ on $\mathbb{R}^d$ and Calderón-Zygmund operators associated to μ on these spaces.

In section 2, we obtain several characterizations in terms of mean oscillations of functions in the Lipschitz spaces associated to μ . Then these characterizations are used in section 3 to establish the boundedness of Calderón-Zygmund operators.

From now on, all balls that we consider will be centered at points in the support of μ .

2. CHARACTERIZATION OF LIPSCHITZ SPACES

In order to prove the main theorem of this section we will need the following known definition and lemma (see [To3])

Definition 2.1. Let β be a fixed constant. A ball B is called β -doubling if

$$\mu(2B) \leq \beta\mu(B).$$

Lemma 2.2. Let $f \in L^1_{\text{loc}}(\mu)$. If $\beta > 2^d$, then, for almost every x with respect to μ , there exists a sequence of β -doubling balls $B_j = B(x, r_j)$ with $r_j \rightarrow 0$, such that

$$\lim_{j \rightarrow \infty} \frac{1}{\mu(B_j)} \int_{B_j} f(y) d\mu(y) = f(x).$$

Proof. We will show that for almost every x with respect to μ there is a β -doubling ball centered at x with radius as small as we wish. This fact, combined with the differentiation theorem, completes the proof of the lemma.

We know that for almost every x with respect to μ

$$(2.1) \quad \lim_{r \rightarrow 0} \frac{\mu(B(x, r))}{r^d} > 0$$

(differentiation of μ with respect to Lebesgue measure, see [M]). Now for x satisfying (2.1) take $B = B(x, r)$ and assume that none of the balls $2^{-k}B$, $k \geq 1$, is β -doubling. Then it is easy to see that $\mu(B) > \beta^k \mu(2^{-k}B)$ for all $k \geq 1$. Therefore

$$\frac{\mu(2^{-k}B)}{(2^{-k}r)^d} < \left(\frac{2^d}{\beta}\right)^k \frac{\mu(B)}{r^d}.$$

Note that, since $\beta > 2^d$, the right hand side tends to zero for $k \rightarrow \infty$, which is a contradiction. $\square$

Now we can state and prove the main result of this section.

Theorem 2.3. *For a function $f \in L^1_{\text{loc}}(\mu)$, the conditions I, II, and III below, are equivalent*

- (I) *There exist some constant C_1 and a collection of numbers f_B , one for each ball B , such that these two properties hold: For any ball B with radius r*

$$(2.2) \quad \frac{1}{\mu(2B)} \int_B |f(x) - f_B| \, d\mu(x) \leq C_1 r^\alpha,$$

and for any ball U such that $B \subset U$ and $\text{radius}(U) \leq 2r$.

$$(2.3) \quad |f_B - f_U| \leq C_1 r^\alpha,$$

- (II) *There is a constant C_2 such that*

$$(2.4) \quad |f(x) - f(y)| \leq C_2 |x - y|^\alpha$$

for μ -almost every x and y in the support of μ .

- (III) *For any given p , $1 \leq p \leq \infty$, there is a constant $C(p)$, such that for every ball B of radius r , we have*

$$(2.5) \quad \left(\frac{1}{\mu(B)} \int_B |f(x) - m_B(f)|^p \, d\mu(x) \right)^{1/p} \leq C(p) r^\alpha,$$

where $m_B(f) = \frac{1}{\mu(B)} \int_B f(y) \, d\mu(y)$ and also for any ball U such that $B \subset U$ and $\text{radius}(U) \leq 2r$.

$$(2.6) \quad |m_B(f) - m_U(f)| \leq C(p) r^\alpha,$$

In addition, the quantities: $\inf C_1$, $\inf C_2$, and $\inf C(p)$ with a fixed p are equivalent.

Proof. (I) $\Rightarrow$ (II). Consider x as in the Lemma and let $B_j = B(x, r_j)$, $j \geq 1$, a sequence of β -doubling balls with $r_j \rightarrow 0$. We will show first that (2.2) implies

$$\lim_{j \rightarrow \infty} f_{B_j} = f(x).$$

It suffices to observe that

$$\begin{aligned} |m_{B_j}(f) - f_{B_j}| &\leq \frac{1}{\mu(B_j)} \int_{B_j} |f(y) - f_{B_j}| \, d\mu(y) \\ &\leq \frac{\mu(2B_j)}{\mu(B_j)} \frac{1}{\mu(2B_j)} \int_{B_j} |f(y) - f_{B_j}| \, d\mu(y) \leq \beta C_1 r_j^\alpha. \end{aligned}$$

Next, let x and y be two points as in the lemma. Take $B = B(x, r)$ any ball with $r \leq |x - y|$ and let $U = B(x, 2|x - y|)$. Now define $B_k = B(x, 2^k r)$, for $0 \leq k \leq \bar{k}$, where $\bar{k}$ is the first integer such that $2^{\bar{k}} r \geq |x - y|$. Then

$$\begin{aligned} |f_B - f_U| &\leq \sum_{k=0}^{\bar{k}-1} |f_{B_k} - f_{B_{k+1}}| + |f_{B_{\bar{k}}} - f_U| \\ &\leq C_1 \sum_{k=0}^{\bar{k}} (2^k r)^\alpha \leq C' C_1 |x - y|^\alpha, \end{aligned}$$

where C' is independent of x and B .

A similar argument can be made for the point y with any ball $B' = B(y, s)$ such that $s \leq |x - y|$ and $V = B(y, 3|x - y|)$. Therefore

$$|f_B - f_{B'}| \leq |f_B - f_U| + |f_U - f_V| + |f_V - f_{B'}| \leq C'' C_1 |x - y|^\alpha.$$

Finally, take two sequences of β -doubling balls $B_j = B(x, r_j)$ and $B'_j = B(y, s_j)$ with $r_j \rightarrow 0$ and $s_j \rightarrow 0$. We have

$$|f(x) - f(y)| = \lim_{j \rightarrow \infty} |f_{B_j} - f_{B'_j}| \leq C'' C_1 |x - y|^\alpha.$$

(II) $\Rightarrow$ (III). It is immediate. Note also that (II) $\Rightarrow$ (I) is immediate as well.

(III) $\Rightarrow$ (I). Define first $f_B = m_B(f)$. Then (2.3) is exactly (2.6).

To prove (2.2), we write

$$\begin{aligned} \frac{1}{\mu(2B)} \int_B |f(y) - f_B| \, d\mu(y) &\leq \frac{1}{\mu(2B)} \left(\int_B |f(y) - f_B|^p \right)^{1/p} \mu(B)^{1/p'} \\ &\leq \frac{\mu(B)}{\mu(2B)} \left(\frac{1}{\mu(B)} \int_B |f(y) - f_B|^p \, d\mu(y) \right)^{1/p} \leq C(p) r^\alpha. \end{aligned}$$

This concludes the proof of the theorem. $\square$

Remark 2.4. Theorem 2.3 is also true if the number 2 in condition (I) is replaced by any fixed $\rho > 1$. In that case, the proof uses (ρ, β) -doubling balls, that is, balls satisfying $\mu(\rho B) \leq \beta \mu(B)$. However this extension is not needed in our paper.

The idea of combining the mean oscillation condition with an extra condition as in (I) originates in the work of Tolsa [To3] on regular BMO , whereas the introduction of the ρ factor in (I) comes from [NTV3].

Definition 2.5. We shall call Lipschitz function of order α with respect to μ to a function, or rather the corresponding Lebesgue class in $L^1_{\text{loc}}(\mu)$, which satisfies any, and hence all, of the conditions of theorem 2.3.

The linear space of all Lipschitz functions of order α , with respect to μ , modulo constants, becomes, with the norm $\inf C_2$ of Theorem 2.3, a Banach space, which we shall call $\mathcal{Lip}(\alpha, \mu)$.

Remark 2.6. It is easy to see that $\mathcal{Lip}(\alpha, \mu)$ coincides with the space of Lipschitz functions of order α on the support of μ .

Note that by the extension theorem of Banach (see [B] or [Mi]), any Lipschitz function of order α with respect to μ has an extension to $\mathbb{R}^d$ that is a Lipschitz function of order α with an equivalent norm.

Remark 2.7. For $0 < \alpha \leq 1$, a telescoping argument like the one used in the proof of (I) $\Rightarrow$ (II) in Theorem 2.3 shows that (2.3) is equivalent to

$$(2.7) \quad |f_B - f_U| \leq C'_1 \text{radius}(U)^\alpha$$

for any two balls $B \subset U$. Furthermore (2.7) is also equivalent to

$$(2.8) \quad |f_B - f_U| \leq C''_1 K_{B,U} \text{radius}(U)^\alpha,$$

for any two balls $B \subset U$, where $K_{B,U}$ is the constant introduced by X. Tolsa in [To3], given by

$$K_{B,U} = 1 + \sum_{j=1}^{N_{B,U}} \frac{\mu(2^j B)}{(2^j B)^n},$$

with $N_{B,U}$ equal to the first integer k such $2^k \text{radius}(B) \geq \text{radius}(U)$. Indeed (2.8) for comparable balls, that is, for $\text{radius}(U) \leq 2 \text{radius}(B)$, reduces to (2.3) because, in this case, $K_{B,U}$ is controlled by an absolute constant.

Note that (2.2) and (2.8) make sense also for $\alpha = 0$ and the space defined by them is the space $RBMO(\mu)$ of X. Tolsa (see [To3]). Therefore, the spaces $\mathcal{Lip}(\alpha, \mu)$, $0 < \alpha \leq 1$ can be seen as members of a family containing also $RBMO(\mu)$.

3. CALDERÓN-ZYGMUND OPERATORS

Definition 3.1. An “ n -dimensional” standard kernel K on $\mathbb{R}^d$ will be a function $K \in L^1_{\text{loc}}(\mathbb{R}^d \times \mathbb{R}^d \setminus \{(x, y) : x = y\}, \mu \times \mu)$, satisfying the following two conditions for some constants $C > 0$ and $\varepsilon \in]0, 1]$:

- (i) $|K(x, y)| \leq \frac{C}{|x - y|^n}$, for all $x \neq y$ and
- (ii) $|K(x, y) - K(x', y)| \leq C \frac{|x - x'|^\varepsilon}{|x - y|^{n+\varepsilon}}$ for $|x - y| \geq 2|x - x'|$.

When we want to be precise about ε , we shall say that K is an n -dimensional, ε -regular, standard kernel.

Definition 3.2. By a Calderón-Zygmund operator on $\mathbb{R}^d$ with respect to μ , we shall mean a linear operator T , bounded on $L^2(\mu)$, which is associated to an n -dimensional standard kernel K in the sense that for every $f \in L^2(\mu)$, with compact support

$$Tf(x) = \int_{\mathbb{R}^d} K(x, y) f(y) d\mu(y), \quad \text{for } \mu\text{-almost every } x \in \mathbb{R}^d \setminus \text{supp } f.$$

Remark 3.3. We have not required ϵ -regularity of K in the second variable, as is usually done, because we do not need it for proving Theorem 3.6.

The definition of a Calderón-Zygmund operator T can be extended to Lipschitz functions in the following way:

Definition 3.4. Assume that T is a Calderón-Zygmund operator associated to μ having an ϵ -regular kernel. First, for each ball B , with center x_B , $x \in B$ and $f \in \mathcal{Lip}(\alpha, \mu)$, $0 < \alpha < \epsilon$, we define

$$(3.1) \quad T_B(f)(x) = T(f\chi_{2B})(x) + \int_{\mathbb{R}^d \setminus 2B} (K(x, z) - K(x_B, z)) f(z) d\mu(z)$$

Now $T(f)$ is defined as the class of all functions g on $\mathbb{R}^d$ such that for any ball B there is a constant c_B satisfying

$$g(x) = T_B(f)(x) + c_B \quad \mu\text{-almost everywhere on } B.$$

Remark 3.5. In order to see that $T(f)$ is well defined, it suffices to realize that if $B \subset V$ are balls, then $T_B(f) - T_V(f)$ is constant μ -a. e. over B . Then, considering an increasing sequence of balls whose union is $\mathbb{R}^d$, we can see that there is a function g such that $g - T_B(f)$ is constant μ -a. e. on B for each ball B in the sequence. Consider now any ball B . Since this ball B is contained in some ball of the sequence, by our previous observation, $g - T_B(f)$ is constant μ -a. e. on B .

Theorem 3.6. Let T be a Calderón-Zygmund operator associated to μ having an ϵ -regular kernel and $0 < \alpha < \epsilon \leq 1$. Then T is bounded on $\mathcal{Lip}(\alpha, \mu)$ if and only if $T(1) = 0$ in $\mathcal{Lip}(\alpha, \mu)$.

Proof. The fact that $T(1) = 0$ in $\mathcal{Lip}(\alpha, \mu)$ is a necessary condition follows immediately from the fact that $\|1\|_{\mathcal{Lip}(\alpha, \mu)} = 0$.

To prove that this condition is sufficient, take $f \in \mathcal{Lip}(\alpha, \mu)$ and two balls $B \subset U$ with $\text{radius}(U) \leq 2r$, where r is the radius of B . We want to show that there are two constants a_B and a_U such that

$$(3.2) \quad \frac{1}{\mu(2B)} \int_B |T_U(f)(x) - a_B| \, d\mu(x) \leq C \|f\|_{\mathcal{Lip}(\alpha, \mu)} r^\alpha$$

and

$$(3.3) \quad |a_B - a_U| \leq C \|f\|_{\mathcal{Lip}(\alpha, \mu)} r^\alpha$$

with C independent of B, U and f .

Since $T(f)(x) = T_U(f)(x) + c_U$ on U , where c_U is a constant, then (3.2) and (3.3) follow for $T(f)$ with constants $a'_B = a_B + c_U$ and $a'_U = a_U + c_U$.

By the hypothesis $T(1) = 0$, we can assume that $\int_{2U} f \, d\mu = 0$. Let's now define

$$a_B = \frac{1}{\mu(2B)} \int_B T_U(f)(x) \, d\mu(x)$$

and, similarly a_U with U instead of B . Observe that

$$\begin{aligned} \frac{1}{\mu(2B)} \int_B |T_U(f)(x) - a_B| \, d\mu(x) &\leq \frac{1}{\mu(2B)} \int_B |T_U(f)(x)| \, d\mu(x) + \frac{1}{\mu(2B)} \int_B |a_B| \, d\mu(x) \\ &\leq 2 \frac{1}{\mu(2B)} \int_B |T_U(f)(x)| \, d\mu(x), \end{aligned}$$

since $\frac{\mu(B)}{\mu(2B)} \leq 1$. On the other hand

$$|a_B - a_U| \leq \frac{1}{\mu(2B)} \int_B |T_U(f)(x)| \, d\mu(x) + \frac{1}{\mu(2U)} \int_U |T_U(f)(x)| \, d\mu(x).$$

Now we want to estimate the quantity

$$(3.4) \quad \frac{1}{\mu(2B)} \int_B |T_U(f)(x)| \, d\mu(x),$$

where B can also be U . To do that we write

$$\begin{aligned} T_U(f)(x) &= T(f\chi_{2B})(x) + T(f\chi_{2U \setminus 2B})(x) \\ &\quad + \int_{\mathbb{R}^d \setminus 2U} (k(x, z) - k(x_U, z)) f(z) \, d\mu(z). \end{aligned}$$

We then have

$$\begin{aligned} & \frac{1}{\mu(2B)} \int_B |T_U(f)(x)| \, d\mu(x) \\ & \leq \frac{1}{\mu(2B)} \int_B |T(f\chi_{2B})(x)| \, d\mu(x) + \frac{1}{\mu(2B)} \int_B |T(f\chi_{2U \setminus 2B})(x)| \, d\mu(x) \\ & + \frac{1}{\mu(2B)} \int_B \left| \int_{\mathbb{R}^d \setminus 2U} (k(x, z) - k(x_U, z)) f(z) \, d\mu(z) \right| \, d\mu(x) = A_1 + A_2 + A_3. \end{aligned}$$

To estimate A_1 , we use that T is bounded in $L^2(\mu)$ and the fact that $|m_{2B}(f)| \leq C \|f\|_{\mathcal{L}ip(\alpha, \mu)} r^\alpha$, which follows from $|m_{2B}(f) - m_{2U}(f)| \leq C \|f\|_{\mathcal{L}ip(\alpha, \mu)} r^\alpha$ and $m_{2U}(f) = 0$. We have

$$\begin{aligned} \frac{1}{\mu(2B)} \int_B |T(f\chi_{2B})(x)| \, d\mu(x) & \leq \left(\int_B |T(f\chi_{2B})(x)|^2 \, d\mu(x) \right)^{1/2} \frac{\mu(B)^{1/2}}{\mu(2B)} \\ & \leq C \left(\int |f\chi_{2B}|^2 \, d\mu \right)^{1/2} \frac{\mu(B)^{1/2}}{\mu(2B)} \\ & \leq C \left(\frac{1}{\mu(2B)} \int_{2B} |f(z) - m_{2B}(f)|^2 \, d\mu(z) \right)^{1/2} \\ & + C \left(\frac{1}{\mu(2B)} \int_{2B} |m_{2B}(f)|^2 \, d\mu(z) \right)^{1/2} \leq C \|f\|_{\mathcal{L}ip(\alpha, \mu)} r^\alpha. \end{aligned}$$

For A_2 , note first that in case $B = U$, A_2 doesn't appear. Otherwise, since $x \notin \text{supp}(f\chi_{2U \setminus 2B})$ and $\mu(2U) \leq C(4r)^n$, we have

$$\begin{aligned} & \frac{1}{\mu(2B)} \int_B \int_{2U \setminus 2B} |k(x, z)| |f(z)| \, d\mu(z) \, d\mu(x) \\ & \leq \frac{1}{\mu(2B)} \int_B \frac{C}{r^n} \int_{2U} |f(z)| \, d\mu(z) \, d\mu(x) \\ & \leq C \frac{\mu(2U)}{r^n} \frac{1}{\mu(2U)} \int_{2U} |f(z) - m_{2U}(f)| \, d\mu(z) \leq C \|f\|_{\mathcal{L}ip(\alpha, \mu)} r^\alpha. \end{aligned}$$

Finally for A_3 using the ϵ -regularity of the kernel, we have

$$\begin{aligned} A_3 & \leq \frac{1}{\mu(2B)} \int_B \int_{\mathbb{R}^d \setminus 2U} \frac{|x - x_U|^\epsilon}{|x_U - z|^{n+\epsilon}} |f(z)| \, d\mu(z) \, d\mu(x) \\ & \leq C \frac{\mu(B)}{\mu(2B)} \|f\|_{\mathcal{L}ip(\alpha, \mu)} r^\alpha \leq C \|f\|_{\mathcal{L}ip(\alpha, \mu)} r^\alpha. \end{aligned}$$

The inequalities above are justified by the fact that, for $z \notin 2U$

$$\begin{aligned} |f(z)| &= |f(z) - m_{2U}(f)| \leq \frac{1}{\mu(2U)} \int_{2U} |f(z) - f(y)| \, d\mu(y) \\ &\leq (2|z - x_U|)^\alpha \|f\|_{\mathcal{L}ip(\alpha, \mu)} \end{aligned}$$

and also, that

$$\int_{\mathbb{R}^d \setminus 2U} \frac{1}{|z - x_U|^{n+\epsilon-\alpha}} \, d\mu(z) \leq Cr^{\alpha-\epsilon}$$

(see [GG]). This concludes the proof of the theorem. $\square$

REFERENCES

- [B] S. Banach *Introduction to the theory of real functions*. Monografie Mat., Tom 17, PWN, Warsaw, 1951.
- [GG] J. García-Cuerva and A. E. Gatto *Boundedness properties of fractional integral operators associated to non-doubling measures*, preprint 2002.
- [MS1] R. A. Macías and C. Segovia *Singular integrals on generalized Lipschitz and Hardy spaces*, Studia Math. 65 (1979) 55–75.
- [MS2] R. A. Macías and C. Segovia *Lipschitz functions on spaces of homogeneous type*, Advances in Math. 33 (1979) 257–270.
- [M] P. Mattila *Geometry of sets and measures in euclidean spaces*. Cambridge studies in advanced Mathematics 44, Cambridge University Press, 1995.
- [Me] Y. Meyer *Ondelettes et opérateurs II. Opérateurs de Calderón-Zygmund*, Hermann, Paris, 1990.
- [MMNO] J. Mateu, P. Mattila, A. Nicolau and J. Orobitg. *BMO for non doubling measures*. Duke Math. J. 102 (2000), no. 3, 533–565.
- [Mi] G. J. Minty *On the extension of Lipschitz, Lipschitz-Hölder continuous and monotone functions*. Bull. Amer. Math. Soc. 76 (1970), 334–339.
- [NTV1] F. Nazarov, S. Treil and A. Volberg. *Cauchy integral and Calderón-Zygmund operators on nonhomogeneous spaces*. Int. Math. Res. Not. 15 (1997), 703–726.
- [NTV2] F. Nazarov, S. Treil and A. Volberg. *Weak type estimates and Cotlar inequalities for Calderón-Zygmund operators in nonhomogeneous spaces*. Int. Math. Res. Not. 9 (1998), 463–487.
- [NTV3] F. Nazarov, S. Treil, and A. Volberg, *Accretive system Tb-theorems on nonhomogeneous spaces*. Duke Math. J. 113 (2002), no. 2, 259–312.
- [S] E. M. Stein *Harmonic analysis. Real-Variable methods, orthogonality, and oscillatory integrals*. Princeton Univ. Press. Princeton, N.J., 1993.
- [To1] X. Tolsa. *L^2 -boundedness of the Cauchy integral operator for continuous measures*. Duke Math. J. 98:2 (1999), 269–304.
- [To2] X. Tolsa. *Cotlar’s inequality and existence of principal values for the Cauchy integral without the doubling condition*. J. Reine Angew. Math. 502 (1998), 199–235.

- [To3] X. Tolsa, *BMO, H^1 and Calderón-Zygmund operators for non-doubling measures*, Math. Ann. 319 (2001), no. 1, 89–149.

DPTO. DE MATEMÁTICAS, C-XV, UNIVERSIDAD AUTÓNOMA, 28049, MADRID, SPAIN

E-mail address: jose.garcia-cuerva@uam.es

DEPARTMENT OF MATHEMATICS, DEPAUL UNIVERSITY, CHICAGO, ILLINOIS, 60614, U. S. A.

E-mail address: aegatto@condor.depaul.edu